



Internationally indexed journal

Indexed in Chemical Abstract Services (USA), Index copernicus, Ulrichs Directory of Periodicals, Google scholar, CABI ,DOAJ , PSOAR, EBSCO , Open J gate , Proquest , SCOPUS , EMBASE ,etc.



Rapid and Easy Publishing

The "International Journal of Pharma and Bio Sciences" (IJPBS) is an international journal in English published quarterly. The aim of IJPBS is to publish peer reviewed research and review articles rapidly without delay in the developing field of pharmaceutical and biological sciences



Pharmaceutical Sciences

- Pharmaceutics
- Novel drug delivery system
- Nanotechnology
- Pharmacology
- Pharmacognosy
- Analytical chemistry
- Pharmacy practice
- Pharmacogenomics
- Polymer sciences
- Biomaterial sciences
- Medicinal chemistry
- Natural chemistry
- Biotechnology
- Pharmaco-informatics
- Biopharmaceutics



Biological Sciences

- Biochemistry
- Biotechnology
- Bioinformatics
- Cell biology
- Microbiology
- Molecular biology
- Neurobiology
- Cytology
- Pathology
- Immunobiology

Indexed in Elsevier Bibliographic Database
(Scopus and EMBASE)
SCImago Journal Rank 0.129
Impact factor 0.47*



*Instruction to Authors visit www.ijpbs.net

For any Queries, visit "contact" of www.ijpbs.net



1-CYSTEINE PEROXIREDOXIN: A LESS STUDIED MEMBER OF PEROXIREDOXIN IN PLANTS

**NEHA GUPTA, MONIKA SHUKLA BAJPAI,
RITA SINGH MAJUMDAR AND PANKAJ K MISHRA***

*Department of Biotechnology, School of Engineering and
Technology, Sharda University, Greater Noida- 201306*

ABSTRACT

1-Cysteine Peroxiredoxin (1Cys Prx) is a thiol dependent antioxidative enzyme of molecular weight 20-30 kDa. It contains one catalytic cysteine residue and is found in cytosol. Prx plays an important role in growth, development and dessication tolerance in dormant seeds. It detoxifies H₂O₂ and various other organic peroxides. It protects lipids, enzymes and DNA against reactive oxygen species. This review highlights our current understanding of 1Cys Prx with focus on its expression in seeds and transgenic plants.

KEYWORDS: Ascorbate Peroxidase, Cysteine, Peroxiredoxin, Glutathione Peroxidase, Hydrogen peroxide, NADPH thioredoxin reductase



PANKAJ K MISHRA

Department of Biotechnology, School of Engineering and
Technology, Sharda University, Greater Noida- 201306

*Corresponding author

INTRODUCTION

Reactive oxygen species (ROS) produced in a cell due to various reasons like incomplete reduction of O_2 to water during respiration, beta-oxidation of fatty acids under stressed conditions exposure to radiation, light, metals & floods. When present in high concentration. They pose an oxidative threat to cells for the scavenging of ROS a number of enzymes are present in the cell which include catalase, glutathione peroxidase, and haem-containing peroxidase & peroxiredoxin. Studies have shown that enhanced levels of ROS lead to redox modifications, such as dithiol-/disulfide transitions, S-nitrosylation, glutathionylation, and subsequently to oxidative damage. A number of ROS are produced in the cell, like superoxide ion (O_2^-), hydroxyl ion (OH^\cdot) and hydrogen peroxide (H_2O_2). Out of these, H_2O_2 is least reactive. H_2O_2 is reduced in the presence of Fe^{2+} in the Fenton reaction and yields hydroxyl radical (OH^\cdot). Due to its ability to diffuse over significant distances within and between cells, H_2O_2 serves a signalling function in cellular communication^{1, 2}. Thus H_2O_2 plays two opposite roles simultaneously, first as a damaging ROS compounds and second as a messenger in cell signalling. For the effective functioning of this molecule its concentration must be balanced in the cell. During normal conditions H_2O_2 concentrations are kept very low. However, level of H_2O_2 increases in response to various abiotic and biotic stresses and takes part in the reactive oxygen regulatory network³. To combat these ROS cells express a number of antioxidative enzymes which decompose H_2O_2 , namely catalase (CAT), ascorbate peroxidase (APX) glutathione peroxidase (GPX), peroxiredoxin (Prx), and type- III peroxidases⁴. Glutathione peroxidase and Prx also decompose alkyl hydroperoxides in addition to H_2O_2 . This review surveys the current literature and presents a comprehensive coverage of current knowledge of 1-Cys Prx with molecular size of 20-30kDa. They are thiol-dependent enzymes containing one (1-cysteine [-Cys]) or two (2-Cys) conserved Cys residues that protect lipids, enzymes, and DNA against ROS. Prx undergoes redox dependent conformational changes and interacts in a conformation-dependent manner with various proteins.

These enzymes are truly ubiquitous having been identified in yeast⁵, plant⁶ and animal cells⁷ including both protozoan and helminth parasites, and most of eubacteria and archaea. The Prxs are primarily located in the cytosol. These (Prx) antioxidants were first identified in yeast thiol-specific antioxidant⁵. Prxs can reduce H_2O_2 , alkyl hydroperoxides, and hydroxyl radicals⁸. *In vitro* they have been shown to have antioxidant activity through protection of biomolecules against radical attack. Based on the sequence, structural similarities and positions of conserved cysteinyl residues the whole Prx family is divided in six groups, named A, B, C, D, E, and F^{9, 10}. Prx from the groups A to D are common and conserved in higher plants. As per the nomenclature, the A-type Prx corresponds to the (typical) 2-Cys Peroxiredoxin (2-CysPrx), the B-type Prx to the (typical) 1-Cys Peroxiredoxin (1-CysPrx), the C-type Prx to peroxiredoxin Q (PrxQ), and the D-type Prx to type II Peroxiredoxins (PrxII). PrxQ and PrxII are also termed atypical 2-Cys Prx¹¹.

Structure and Mode of Action of 1 Cys Peroxiredoxin


1 Cys Peroxiredoxin in plants is one of the very less studied member of Peroxiredoxins. The crystal structure of this enzyme reveals that it exists in a head to tail dimeric structure as shown in (Fig1). This 25 kDa protein contains a single conserved catalytic cysteine (47th position) towards the NH_2 terminal and is thus known as 1-Cys Prx¹². The peroxidase reaction is composed of two steps which are centered on this redox-active cysteine also known as the peroxidatic cysteine. It has been found that site directed mutagenesis of this Cysteine to Serine removes the peroxidase activity of the enzyme^{13, 14, 15}. A number of recent reviews have addressed the structural and catalytic properties of Prx isoforms in general¹⁶ and in plants specifically^{17, 18, 19}. The crystal structure of 1-Cys Prx shows the peroxidatic cysteine as a sulphenic acid ($-SOH$). This sulphenic acid is highly stable in 1-Cys Prx. Close to this peroxidatic cysteine residue His 39 and the Arg 132 is present. This histidine is conserved for only this class

of Prxs enzymes. The presence of these amino acids stabilizes the Cys 47 thiolate anion. 1-Cys Prx act by reducing peroxide substrates to the corresponding alcohol and water. The reaction mechanism comprises of three steps:

1. Oxidation,
2. Derivatization and
3. Regeneration of ground state

In the first catalytic cycle, the peroxidatic thiol group is oxidized to the sulphenic acid and the reduced product, i.e. H₂O in case of H₂O₂, alcohol in case of alkyl hydroperoxides and nitrite in case of alkoxy nitrites is released. The sulphur atom is covered by sulfenate oxygen to prevent its over oxidation to inactive sulfenic acid (SO₂H) and sulfenic acids (SO₃H). There are reports of over oxidation of Prxs¹⁶ as shown in (Fig 2). It is possible due to local unfolding of the active site which exposes the Cys to accomplish this. The structures have shown that C terminal arm gets unfolded, positioning the resolving Cys for disulphide bond formation. This disulphide bond is exposed at one end, thereby allowing the resolving cysteine to be attacked by thiol

containing reductants. Pulido *et al*²⁰ have shown that the NADPH thioredoxin reductase (NTR)-mediates the regeneration of 1-Cys Prx in the nuclei of wheat seed cells exposed to various oxidative stresses. Crystal structures of the disulfide-bonded enzymes have shown that the C-terminal arm is unfolded beyond the resolving cysteine, indicating a high degree of mobility for this locally unfolded segment. The overall reaction can be summed as:



1 Cys Prx is a non heme containing enzyme; so it relies on an external electron donor to compensate for the lack of the prosthetic group. The oxidized Prx lose their enzymatic activity and must first be regenerated prior to the next catalytic cycle. For many Prx, it has been established that Thioredoxin serve as electron donor for the regeneration of the active form²¹. It has been shown that the yeast mitochondrial 1-Cys Prx is reactivated by glutathionylation of the catalytic cysteine residue and reduced by thioredoxin reductase coupled with glutathione. Prx 1 plays an important role in protection against mitochondrial oxidative stress²².




Figure 1
Structure of 1 Cys Prx as generated by using MOLSCRIPT and TOPS.




Figure 2
Catalytic mechanism of 1- Cys Prx in peroxide detoxification.

1 Cys Peroxiredoxin in Seeds /Plants

ROS are the molecules which intervene in cellular signalling and growth processes occurring at early embryogenesis and gene expression during seed development²³. Studies have shown that reactive oxygen species are involved in various aspects of programmed cell death²⁴, seed physiology and desiccation²⁵, seed germination and dormancy²⁶, ageing, cellular damage. In seeds, 1 Cys Prx plays an important role in combating the oxidative stress and help in the survival of seeds of the plants. Detoxification done by 1 Cys Prx in vital acquisition of desiccation tolerance of developing seeds, their germination and storability. 1-Cys Prx genes are expressed solely in seeds, and only in the parts of the seeds surviving desiccation, i.e. the embryo and the aleurone layer. 1-Cys Prx found in seeds have been characterized in several species, examined at both the transcript and protein level in relation to dormancy, and described in relation to function and localization. 1-Cys Prx have been identified in a number of plants at gene level like pBS128 in brome grass. Per1 in barley (also called B15C) and FePer1 in buckwheat (*Fagopyrum esculentum*)²⁷ are expressed in developing seeds. The studies have shown that the PER1 the first antioxidant belonging to the 1-Cys subgroup protein of barley is present in high concentrations in the nucleus at the onset of desiccation. Barley seed Peroxiredoxin is encoded by a single Per 1 gene. The studies have shown that levels of 1-Cys transcript can be up regulated by ABA and osmotic stresses and can be suppressed by gibberellic acid. The dormancy-related Prx antioxidant, PER1, is localized in the nucleus of barley embryo and aleurone cells and the protein level is maintained in imbibed dormant seeds. In contrast to most seed-expressed

antioxidants Per1 disappears in germinating embryos, and in the mature aleurone, the transcript is down-regulated by the germinating embryo or by gibberellic acid (GA)²⁸. Dominguez *et al.* studied the nuclear NADPH thioredoxin reductase NTR/1-Cys Prx system and found that NADPH is able to detoxify H₂O₂ under oxidative stress conditions in cereal seeds²⁹. It controls the stress conditions in nucleus which is important for redox regulation of gene expression in germinating seed cells. The studies have shown NADPH thioredoxin reductase (NTR) supports the antioxidant activity of 1-Cys Prx as shown in (Fig 3). This Redox system is also localized in the nucleus of wheat seed cells and probably plays an important role in antioxidant protection in aleurone and scutellum cells suffering from oxidative stress during seed development and germination. The pattern of localization of NADPH thioredoxin reductase NTR in developing and germinating wheat seeds were studied using an immunocytochemical analysis. The presence of NTR in transfer cells, vascular tissue, developing embryo and root meristematic cells, agrees with the localization of Trx h, and it supports the important function of the NTR/Trx system in cell proliferation and communication. Interestingly, NTR is found to be co-localized with Trx and 1 Cys Prx in the nuclei of seed cells suffering oxidative stress²⁰. A full length cDNA encoding 1 Cys Prx was cloned from wheat and the recombinant protein was expressed in *Escherichia coli*, to test whether the NTR/Trx systems serves as an electron donor. From the purified components, it was found that activity of 1 Cys Prx is NTR dependent. Mutants of the 1-Cys Prx show that the peroxidatic residue of the wheat enzyme is Cys46, which is over oxidized *in vitro* under oxidant conditions. The

extracts from developing and germinating seeds were analysed and it was found that there is over oxidation of 1-Cys Prx *in vivo*. Based on these results, it was proposed that NADPH is the source of the reducing power to regenerate 1-Cys Prx in the nuclei of wheat seeds cells suffering oxidative stress, in a process that is catalyzed by NTR³⁰. 1-Cys Prx gene has also been identified in a medicinal Mushroom *Antrodia camphorata*³¹. It was found to contain 23 amino acid residues with calculated molecular mass of 25,081 Da. The protein obtained shared 44-58% identity with 1-Cys Prx from *Homo sapiens*, *Bos Taurus* and *Saccharomyces cerevisia*. The purified recombinant enzyme was thermo stable with a half life of inactivation 15.5 min at 60degree C. It was stable in the pH range 7.8-10.2. Ikegami in (2009)³² showed that 1 Cys

Prx (DkPrx) is present in abundance in fruits of persimmon (*Diospyros kaki* Thunb.) which accumulate large amounts of proanthocyanidins (PAs) in early stages of development. The DkPrx reported is the first member of 1Cys Prx reported from fruits of any plant species. However, whether this enzyme play role in biosynthesis/accumulation of PAs or it is the accumulation of PAs that induce its gene expression is not clear yet. Peroxiredoxins are one of the key players of the thiosulfide redox regulatory network of plant cells. The role of these peroxiredoxins was studied in *Lotus japonicas*³³ and it was found that 1 Cys Prx gene shows high expression in the seed embryos and low expression in vegetative tissues. The expression was induced by nitric oxide and cytokinin.




Figure 3

NTR and 1 Cys Prx forms nuclear localized redox system under oxidative stress.

Chaperone Activity of 1 Cys Peroxiredoxin

Over oxidation of the conserved cys residue, on H₂O₂ treatment increased its chaperone activity two folds. However, its peroxidase activity decreased considerably without enzyme. Kim et al. (2011)³⁴ have shown that during seed germination 1-Cys Prx not only relieve mild oxidative stress but also act as molecular chaperons and it is the overoxidation that controls the switch in function of 1-Cys Prx from Peroxidase to molecular chaperones.

1 Cys Peroxiredoxins in Transgenic Plants

Proteomic analysis of rice callus was done and 10 ABA induced proteins were identified as the product of the embryo specific promoter

candidates. 1 Cys Prx, was cloned and analyzed. Beta glucuronidase (GUS) expression driven by 1 Cys Prx promoters increases by ABA treatment and rapidly induced by wounding in callus and at the leaf of the transgenic plants. Ectopic expression of the GUS construct in *Arabidopsis* suggested that the 1-Cys Prx promoter also has strong activity in seeds of dicot plants^{35, 36}. The transcript level of rice 1Cys-Prx (R1C-Prx) rapidly decreased after imbibition of rice seeds, but the protein was detected for 15 days after imbibitions. Transgenic tobacco plants constitutively expressing the R1C-Prx gene were considered to study the function of 1 Cys Prx. These plants showed a germination frequency similar to control

plants, however, they exhibited higher resistance against oxidative stress, suggesting that antioxidant activity may be the primary function of 1 Cys Prx³⁷. 1-Cys Prx has been shown to be higher in seedlings of soybean (*Glycine max* (L.) Merr.) suffering from flooding stress than in normally growing seedlings. In a study, soybean genome was found to contain two genes corresponding to 1-Cys Prx and is designated as GmPer1a, GmPer1b. GmPer1a encodes a polypeptide containing the putative catalytic site, and the GmPer1b contains a stop codon inside the deduced polypeptide-coding region, indicating that GmPer1b might be a pseudogene. Two gene forms of GmPer1 protein existed in both submerged and normally grown seedlings, and both forms were higher in the submerged seedlings³⁸. From *Xerophyta viscosa* a monocotyledonous desiccation tolerant plant, a stress inducible gene named XvPer1 has been isolated. The length of the cDNA sequence was found to be 849bp, containing one major reading frame of 660 bp. It was found to exhibit 70% similarity with other 1 Cys Prx proteins. The amino acid sequence constituting the active site of the enzyme, PVCTTE, has been found to be highly conserved in XvPer1. Western blot analysis

and immunofluorescence analysis revealed that XvPer1 is localized in the nucleus of dehydrated *Xerophyta viscosa* leaf cells. These results suggest that XvPer1 is a stress-inducible gene, which may function to protect nucleic acids within the nucleus against oxidative injury¹².

CONCLUSION

1 Cys Prx is a ubiquitous enzyme playing diverse roles ranging from the antioxidant defense of the cell³⁹ to growth, development and desiccation tolerance in dormant seeds. They are known to catalyze a diverse range of peroxides by using a thiol based catalytic mechanism. It efficiently works at low peroxide concentrations. This enzyme is also involved in redox dependent signalling by altering H₂O₂ concentration, processing alkyl hydroperoxides and decomposing peroxyxynitrite. They perform redox interactions with other proteins and undergo redox dependent conformational changes. The literature within the 1 Cys prx field is currently focussed on their more recently identified roles as molecular chaperones under oxidative stress conditions.

REFERENCES

1. Foyer H and Noctor G, Oxygen processing in photosynthesis: regulation and signaling, *New Phytol.*, 146: 359–388, (2000).
2. Apel K and Hirt H, Reactive oxygen species: metabolism, oxidative stress and signal transduction, *Ann. Rev. Plant Biol.*, 55: 373–399, (2004).
3. Mittler R, Vanderauwera S, Gollery M and Van Breusegem F, Reactive oxygen gene network of plants. *Trends Plant Sci.*, 9: 490–498, (2004).
4. Mittler R and Poulos TL, Ascorbate peroxidase. In: Smirnoff N, ed. *Antioxidants and reactive oxygen species in plants*, Oxford: Blackwell Publishing, 87–100, (2005).
5. Nicolas Rouhier, Eric Gelhaye, Pierre-Eric Sautiere, Annick Brun, Pascal Laurent, Denis Tagu, Joelle Gerard, Elisabeth de Fay, Yves Meyer, and Jean-Pierre Jacquot, Isolation and Characterization of a New Peroxiredoxin from Poplar Sieve Tubes That Uses Either Glutaredoxin or Thioredoxin as a Proton Donor. *Plant Physiol.*, 127: 1299–1309, (2001).
6. Dietz KJ, Peroxiredoxins in Plants and Cyanobacteria, *Antioxidants & redox signaling*, 15: 1130-1152, (2011).
7. Kim HS, Manevich Y, Feinstein SI, Pak JH, Ho YS and Fisher AB, Induction of 1-Cys Peroxiredoxin expression by oxidative stress in lung epithelial cells. *Am. J. physiol. lung cell mol. physiol.*, 285: 363-369, (2003).
8. Gao C, Zhang K, Yang G and Wang Y, Expression Analysis of Four Peroxiredoxin Genes from *Tamarix hispida* in Response to Different Abiotic Stresses and Exogenous Abscisic Acid

- (ABA), *Int. J. Mol. Sci.*, 13: 3751-3764, (2012).
9. Noguera-Mazon V, Lemoine J, Walker O, Rouhier N, Salvador A, Jacquot JP, Lancelin JM and Krimm I, Glutathionylation induces the dissociation of 1-Cys Peroxiredoxin non-covalent homodimer, *J. Biol. Chem.*, 281: 31736–31742, (2006).
 10. Noh YH, Baek JY, Jeong W, Rhee SG and Chang TS, Sulfiredoxin translocation into mitochondria play a crucial role in reducing hyperoxidized peroxiredoxin III, *J. Biol. Chem.*, 284: 8470–8477, (2009).
 11. Hofmann B, Hecht HJ and Flohe L, Peroxiredoxins. *Biol. Chem.*, 3833: 47–64, (2002).
 12. Mowla SB, Thomson JA, Farrant JM and Mundree SG, A novel stress-inducible antioxidant enzyme identified from the resurrection plant *Xerophyta viscosa* Baker, *Planta*, 215: 716–726, (2002).
 13. Konig J, Baier M, Horling F, Kahmann U, Harris G, Schurmann P and Dietz KJ, The plant-specific functions of 2-Cys peroxiredoxin-mediated detoxification of peroxides in the redox-hierarchy of photosynthetic electron flux, *Proc. Natl. Acad. Sci. USA.*, 99: 5738–5743, (2002).
 14. Rouhier N, Gelhaye E, Gualberto JM, Jordy MN, De Fay E, Hirasawa M, Duplessis S, Lemaire SD, Frey P, Martin F, Manieri W, Knaff DB and Jacquot JP, Poplar peroxiredoxin Q. A thioredoxin-linked chloroplast antioxidant functional in pathogen defense, *Plant Physiol.*, 134: 1027–1038, (2004).
 15. Lamkemeyer P, Laxa M, Collin V, Li W, Finkemeier I, Schottler MA, Holtkamp V, Tognetti VB, Issakidis-Bourguet E, Kandlbinder A, Weis E, Miginiac-Maslow M and Dietz KJ, Peroxiredoxin Q of *Arabidopsis thaliana* is attached to the thylakoids and functions in context of photosynthesis, *Plant J.*, 45: 968–981, (2006).
 16. Wood ZA, Schroder E, Harris JR and Poole LB, Structure, mechanism and regulation of peroxiredoxins, *TRENDS in Biochemical Sciences*, 28: 32-40, (2003).
 17. Dietz KJ, Horling F, Konig J and Baier M, The function of the chloroplast 2cysteine peroxiredoxin in peroxide detoxification and its regulation, *J. Exp. Bot.*, 53: 1321–1329, (2002).
 18. Rouhier N, Gelhaye E and Jacquot JP, Glutaredoxin-dependent peroxiredoxin from poplar: protein-protein interaction and catalytic mechanism, *J. Biol. Chem.*, 277: 13609–13614, (2002).
 19. Dietz KJ, *Plant Peroxiredoxins*, *Annu. Rev. Plant Biol.*, 54: 93–107, (2003).
 20. Pulido P, Cazalis R and Cejudo FJ, An antioxidant redox system in the nucleus of wheat seed cells suffering oxidative stress, *Plant J.*, 57: 132–145, (2009).
 21. Dietz KJ, Jacob S, Oelze ML, Laxa M, Tognetti V, Miranda SM De, Baier M and Finkemeier I, The function of peroxiredoxins in plant organelle redox metabolism, *J. Exp. Bot.*, 57: 1697–1709, (2004).
 22. Greetham D and Grant CM, Antioxidant Activity of the Yeast Mitochondrial One-Cys Peroxiredoxin Is Dependent on Thioredoxin Reductase and Glutathione In Vivo, *Mol. and Cellu. Biol.*, 3229–3240, (2009).
 23. Bailly C, Active oxygen species and antioxidants in seed biology, *Seed Sci. Res.*, 14: 93-107, (2004).
 24. Fath A, Bethke P, Beligni V and Jones R, Active oxygen and cell death in cereal aleurone cells. *J. Exp. Bot.*, 53: 1273-1282, (2002).
 25. Leprince O and Buitink J, Desiccation tolerance: From genomics to the field, *Plant Sci.*, 179: 554–564, (2010).
 26. Penfield S and King J, Towards a systems biology approach to understanding seed dormancy and germination, *Proc. R. Soc. B.*, 276: 3561–3569, (2009).
 27. Lewis ML, Miki K and Ueda T, FePer 1, a gene encoding an evolutionarily conserved 1-Cys peroxiredoxin in buckwheat (*Fagopyrum esculentum* Moench), is expressed in a seed-specific manner and induced during seed germination, *Gene*, 246: 81–91, (2008).
 28. Serrato AJ, Perez-Ruiz JM and Cejudo FJ, Cloning of thioredoxin reductase and characterization of the thioredoxin reductase-thioredoxin system from wheat, *Biochem. J.*, 217: 392–399, (2002).

29. Domínguez F, Moreno J and Cejudo FJ, A gibberellin-induced nuclease is localized in the nucleus of wheat aleurone cells undergoing programmed cell death, *J. Biol. Chem.*, 279: 11530-6, (2004).
30. Pulido P, Dominguez F and Cejudo FJ, A hydrogen peroxide detoxification system in the nucleus of wheat seed cells, *Plant Signalling and Behavior.*, 4: 23-25, (2009).
31. Wen L, Huang HM, Juang RH and Lin CT, Biochemical characterizations of 1-Cys peroxiredoxin from *Antrodia camphorate*, *Appl. Microbiol. Biotechnol.*, 73: 1314-1322, (2007).
32. Ikegami A, Akagi T, Potter D, Yamada, M, Sato A, Yonemori K, Kitajima A and Inoue K, Molecular identification of 1-Cys peroxiredoxin and anthocyanidin/Xavonol 3-O-galactosyltransferase from proanthocyanidin-rich young fruits of persimmon (*Diospyros kaki* Thunb.), *Planta*, 230: 841–855, (2009).
33. Mendez AT, Matamoros MA, Sanmamed PB, Dietz KJ, Cejudo FJ, Rouhier N, Sato S, Tabata S and Becana M, Peroxiredoxins and NADPH-Dependent Thioredoxin Systems in the Model Legume *Lotus japonicas*, *Plant Physiol.*, 156: 1535–1547, (2011).
34. Kim JH, Jung IJ, Kim DY, Fanata WI, Son BH, Yoo JY, Harmoko R, Ko KS, Moon JC, Jang HH, Kim WY, Kim JY, Lim CO, Lee SY and Lee KO, Proteomic identification of an embryo-specific 1Cys-Prx promoter and analysis of its activity in transgenic rice, *Biochemical and Biophysical Research Communications*, 408: 78-83, (2011).
35. Kim SY, Paeng SK, Nawkar GM, Maibam P, Lee ES, Kim KS, Lee DH, Park DJ, Kang SB, Kim MR, Lee JH, Kim YH, Kim WY and Kang CH, The 1-cys peroxiredoxin, a regulator of seed dormancy, functions as a molecular chaperone under oxidative stress conditions, *Plant Sci.*, 181: 119-24, (2011).
36. Haslekas C, Grini PE, Nordgard SH, Thorstensen T, Viken MK, Nygaard V and Aalen RB, ABI3 mediates expression of the peroxiredoxin antioxidant AtPER1 gene and induction by oxidative stress. *Plant Mol. Biol.*, 53: 313–326, (2003).
37. Lee KO, Jang HH, Jung BG, Chi YH, Lee JY, Choi YO, Lee JR, Lim CO, Cho MJ and Lee SY, Rice 1 Cys-peroxiredoxin over-expressed in transgenic tobacco does not maintain dormancy but enhances antioxidant activity, *FEBS Lett.*, 486: 103-106, (2000).
38. Nishizawa K and Komatsu S, Characteristics of soybean 1-Cys peroxiredoxin and its behavior in seedlings under flooding stress. *Plant Biotech.*, 28: 83–88, (2011).
39. Rabilloud T, Heller M, Gasnier F, Luche S, Rey C, Aebersold R, Mohamed Benahmed, Louisot P and Lunardi J, Proteomics analysis of cellular response to oxidative stress: evidence for in vivo over-oxidation of peroxiredoxins at their active site, *J. Biol. Chem.*, 277: 19396–19401, (2002).